

AD-A206 152

REPORT DOCUMENTATION PAGE

(1)

Unclassified			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NMRI 89-14			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION Naval Medical Research		6b. OFFICE SYMBOL (if applicable)	7a. NAME OF MONITORING ORGANIZATION Naval Medical Command	
6c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814-5055			7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Washington, D.C. 20372-5120	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Naval Medical Research and Development Command		8b. OFFICE SYMBOL (if applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Bethesda, Maryland 20814-5055			10. SOURCE OF FUNDING NUMBERS	
			PROGRAM ELEMENT NO. 62758N	PROJECT NO. MF58524
			TASK NO. 02C-0007	WORK UNIT ACCESSION NO. DN077674
11. TITLE (Include Security Classification) Intensity Threshold for 60-Hz Magnetically Induced Behavioral Changes in Rats				
12. PERSONAL AUTHOR(S) Liboff AR, Thomas JR, Schrot J				
13a. TYPE OF REPORT journal article		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 1989
15. PAGE COUNT 3				
16. SUPPLEMENTARY NOTATION Reprinted from: Bioelectromagnetics 1989; Vol.10 pp. 111-113				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	cyclotron resonance DRL schedule	
			timing discrimination	
			A.C. threshold	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)				
DTIC SELECTED MAR 27 1989 DCE				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED / UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Phyllis Blum, Information Services Division			22b. TELEPHONE (include Area Code) 202-295-2188	22c. OFFICE SYMBOL ISD / ADMIN / NMRI

Brief Communication

Intensity Threshold for 60-Hz Magnetically Induced Behavioral Changes in Rats

Abraham R. Liboff, John R. Thomas, and John Schrot

Department of Physics, Oakland University, Rochester, Michigan (A.R.L.); Behavior
Technology Branch, Environmental Stress Program Center, Naval Medical Research
Institute, Bethesda, Maryland (J.R.T., J.S.)

Experiments were conducted to further investigate the effect of 60-Hz cyclotron-resonance exposures on rats performing on a multiple FR-DRL schedule. The previously reported temporary loss of DRL baseline response, when measured as a function of A.C. magnetic intensity, was found to have a threshold. Utilizing the component of A.C. magnetic intensity parallel to the D.C. field, we report this threshold as $(0.27 \pm 0.10) \times 10^{-4} T_{rms}$.

Key words: cyclotron resonance, timing discrimination, A.C. threshold, DRL schedule

Received for publication 10/15/87; accepted 1/15/88; reprint requests to (A.R.L.)

INTRODUCTION

We recently reported [Thomas et al., 1986] that rats operantly conditioned on a multiple, fixed ratio differential low-rate (FR-DRL) schedule exhibited a temporary loss of their DRL-baselines following a 30-min exposure to a specific combination of magnetostatic ($0.261 \times 10^{-4} T$) and 60-Hz linearly polarized magnetic fields. The multiple schedule was such that when a red light was on the rat performed on an FR or counting schedule, pressing a lever 30 times (FR 30) to produce a food pellet. When a green light was illuminated a DRL schedule was in effect, with a pellet produced by a response on the lever following a preceding lever response by at least 18 s (DRL 18). Although performance on the FR schedule was not systematically affected by field exposure, there was a consistent modification in both the rate and pattern of responding on the DRL schedule following exposure to a combination of magnetic fields chosen to correspond to the cyclotron resonance tuning condition for

Received for review June 15, 1987; revision received July 15, 1988.

Address reprint requests to A. R. Liboff, Department of Physics, Oakland University, Rochester, MI 48309-4401.

1989 Alan R. Liss, Inc.

89 3 24 030

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20






Fig. 1. Field configuration used by Thomas et al. [1986]. A: Magnetostatic components: $B_V = 0.157 \times 10^{-4}$ T, $B_H = 0.209 \times 10^{-4}$ T, $B_T = 0.261 \times 10^{-4}$ T. Direction B_H corresponds to compass heading (local geomagnetic north). In addition, 60-Hz linearly polarized magnetic field B_{AC} applied parallel to B_H . B: Construction used to obtain B_{AC} , the component of B'_{AC} in the direction of B_T . It follows that $B_{AC} = B'_{AC} \sin \theta = 0.801 B'_{AC}$.

the singly charged lithium ion. The 60-Hz magnetic field exposure was directed horizontally within the N-S geomagnetic plane along the long axis of the rat. The minimum intensity of this field was 5×10^{-5} T_{rms}. The magnetostatic field was adjusted by partial compensation of the local geomagnetic field to a net intensity of 0.261×10^{-4} T at an angle of 36.8° to the horizontal (see Fig. 1).

It is important to point out that the cyclotron resonance condition that we applied in the original experiment does not depend on the intensity of the A.C. field. This condition is solely dependent on three variables: the frequency of the A.C. field, the magnetostatic intensity, and the ionic charge-to-mass ratio—in the present case, respectively, 60 Hz, 0.261×10^{-4} T, and 1.39×10^7 Coul/kg. The role of the A.C. magnetic intensity in the interaction mechanism is not clear, except that it represents a potential means (albeit ultraweak) of energy transfer. The A.C. magnetic intensity also has practical significance since it is used as an exposure parameter in studying those biological effects resulting from power lines and other electromagnetic sources. It is therefore of some interest to determine how the effect reported by Thomas et al. [1986] varies with A.C. magnetic intensity under constant cyclotron resonance conditions.

Additional trials with the same animals were therefore conducted. Except for the range of A.C. magnetic intensities, these additional runs were carried out in precisely the same way as described in the initial report. The present results indicate that the loss of DRL baselines only occurs for 60-Hz intensities in excess of a clearly defined threshold.

The specific configuration of fields that were employed in both the initial and later experiments is shown in Figure 1. Note that B_{AC} , the component of the 60-Hz field that is parallel to the total magnetic field vector (B_T), is only 80% of the actual applied field, B'_{AC} . The values of field that were selected for the initial experiment were chosen by using the fact that 0.261×10^{-4} T is close to cyclotron resonance for lithium ions at 60 Hz. Therefore, in examining the dependence of behavioral response upon A.C. magnetic intensity, we restrict our consideration to that component of the A.C. signal (i.e., B_{AC}) that is in cyclotron resonance with the magnetostatic field B_T .

In Figure 2 the relative (compared to sham) response rate, averaged over the five individual mean response rates following 30-min exposures to 0.261×10^{-4} T, 60 Hz, is plotted as a function of the rms intensity of the A.C. field. The solid curve




Fig. 2. Variation in the relative response rate as a function of rms intensity of 60-Hz magnetic field. Each of the six points shown represents the mean value averaged over the same five rats used in Thomas et al. [1986]. The coordinates of these points correspond to the applied field intensity B'_{AC} . The dashed curve indicates how this curve is shifted toward lower values when determining the response as a function of B_{AC} . By assuming that the mean level for this transition occurs halfway between the sham response level and the higher response plateau, we estimate the threshold intensity for this transition at $0.27 \times 10^{-4} T_{rms}$.

represents the variation with B'_{AC} and the dashed curve represents this variation as a function of B_{AC} . We estimate that the mean threshold for loss of DRL response rate is $B_{AC} = 0.27 \times 10^{-4} T_{rms}$. Since the uncertainty in this value is dominated by the separation of the two points ($0.2 \times 10^{-4} T$) just below and just above the transition region, we report the threshold intensity for the behavioral effect previously reported as $(0.27 \pm 0.10) \times 10^{-4} T_{rms}$.

This study was part of a research program to determine the possible adverse health effects resulting from exposure to the electric and magnetic fields of overhead high-voltage transmission lines, administered by the State of New York Department of Health and Health Research, Inc. The study was also supported by the U.S. Naval Medical Research and Development Command Work Unit No. MF58.524.02C.0001.

REFERENCES

- Thomas JR, Schrot J, Liboff AR (1986): Low-intensity magnetic fields alter operant behavior in rats. *Bioelectromagnetics* 7:349-357.